

Palassini and Young reply: In recent work [1], we studied numerically the ground state structure of the nearest-neighbor Ising spin glass with Gaussian couplings in dimensions, $d = 3, 4$. We looked for large-scale, low-energy excitations about the ground state in finite systems of size $N = L^d$ with periodic boundary conditions, by applying a bulk term in the Hamiltonian which couples to the ground state. The results suggest that there are excitations which flip a finite fraction of spins, have energy which varies as $L^{\theta'}$ with $\theta' \simeq 0$, and have a surface with fractal dimension d_s less than d . The exponent θ' seems to be different from the exponent θ which determines the energy difference when the boundary conditions are changed, e.g. from periodic to anti-periodic, since $\theta > 0$ in both $d = 3, 4$. Similar results were obtained by Krzakala and Martin [2] using a different technique.

In their comment on Ref. [1], Newman and Stein [3] (referred to as NS) have clarified the nature of these excitations (assuming they persist in the thermodynamic limit). They consider the problem from the point of view of “thermodynamic states”, defined in a precise way following earlier work by Fisher and Huse [4] and themselves [5]. In this approach one considers a system of size L_B to be embedded in a much larger system of size L . Roughly speaking, different thermodynamic states correspond to different sets of correlation functions in the size- L_B region as $L \rightarrow \infty$. “Extremal” thermodynamic states are called “pure” states. NS argue that system-size excitations with $\theta' = 0$ and $d_s < d$ would not give rise to new pure states because such excitations would “deflect to infinity” rather than run through the size- L_B system. Only if $d_s = d$ would system-size excitations with $\theta' = 0$ give rise to new thermodynamic states.

The conclusions of NS are consistent with our results, since we did not claim that these excitations would give new pure states. Such excitations could however, as NS also note, be relevant for the excitation spectrum of spin glasses. They could also give rise to a non-trivial order parameter function $P(q)$ on the scale of the whole system, as has been seen in Monte Carlo simulations down to very low temperatures [6].

Of course, another important issue is whether our results, which emerge from studies of small systems, con-

tinue to be valid for large sizes. Based on studies of very large systems in two dimensions, where however $T_c = 0$, Middleton [7] has recently suggested that finite-size corrections may be large in our data because we average over fluctuations of different length scales. Middleton also proposes that our results may go over to those of the “droplet picture” [4] in which $\theta' = \theta (> 0)$ (and $d_s < d$). Our results have also been criticized in the opposite sense by Marinari and Parisi [8] who agree with $\theta' = 0 (\neq \theta)$, but argue, from a different analysis and using their own data, that $d_s = d$, which gives the “replica symmetry breaking” picture.

To be sure of the correct behavior in the thermodynamic limit may require the study of sizes much larger than the sizes currently accessible to simulations. However, an important question which can potentially be addressed now is why there are two different (possibly effective) exponents, θ' and θ , characterizing large-scale excitations of different geometry for the *same* range of sizes.

We thank D. Stein and C. Newman for a useful correspondence.

Matteo Palassini¹ and A.P. Young²,

¹University of California San Francisco

²University of California Santa Cruz

-
- [1] M. Palassini and A. P. Young, Phys. Rev. Lett. **85**, 3017 (2000).
 - [2] F. Krzakala and O. C. Martin, Phys. Rev. Lett. **85**, 3013 (2000).
 - [3] C. M. Newman and D. L. Stein, (unpublished), referred to as NS.
 - [4] D. S. Fisher and D. A. Huse, Phys. Rev. B **38**, 386 (1988).
 - [5] See e.g. C. M. Newman and D. L. Stein, Phys. Rev. E **55**, 5194 (1997).
 - [6] H. Katzgraber, M. Palassini and A. P. Young, cond-mat/0007113.
 - [7] A. A. Middleton, cond-mat/0007375
 - [8] E. Marinari and G. Parisi, cond-mat/0007493